DENNISON, SCHEINER, SCHULTZ & WAKEMAN 612 CRYSTAL SQUARE 4 LAW OFFICES

1745 JEFFERSON DAVIS HIGHWAY ARLINGTON, VIRGINIA 22202-3417

703 412-1155

REMARKS

The claims have been amended to better define the present invention.

Respectfully submitted,

Ira J. Schultz Registration No. 28666

APPENDIX

- 1. (Twice Amended) A process for the preparation of acetic acid, methyl acetate or both acetic acid and methyl acetate, [in a liquid phase reaction medium] comprising a reaction selected from the group consisting of carbonylation of methanol, isomerization of methyl formate or both [processes] carbonylation of methanol and isomerization of methyl formate, said reaction being carried out in a liquid phase reaction medium in the presence of water, a solvent, a homogeneous catalyst system comprising iridium and a halogen-containing promoter, and carbon monoxide, wherein said catalyst system also comprises platinum.
- 2. (Twice Amended) The process as claimed in claim 1 [which is a process for] wherein said reaction is the carbonylation of methanol wherein a carbon monoxide partial pressure of between $0.1.10^5$ Pa and 200.10^5 Pa is maintained throughout the [process] reaction.
- 3. (Twice Amended) The process as claimed in claim 1 [which is a process for] wherein said reaction is the isomerization of methyl formate wherein a carbon monoxide partial pressure of between $0.1.10^5$ Pa and 25.10^5 Pa is maintained throughout the [process] reaction.
- 4.(Twice Amended) The process as claimed in claim 1 [which] wherein said reaction comprises a simultaneous

LAW OFFICES
DENNISON, SCHEINER, SCHULTZ & WAKEMAN
612 CRYSTAL SQUARE 4

1745 JEFFERSON DAVIS HIGHWAY ARLINGTON, VIRGINIA 22202-3417 methanol carbonylation reaction and a methyl formate isomerization reaction, wherein said [process] $\underline{\text{reaction}}$ is carried out under a carbon monoxide partial pressure of between $0.1.10^5$ Pa and 25.10^5 Pa throughout the [process] $\underline{\text{reaction}}$.

30.(Amended) The process as claimed in claim 21 which is carried out in the presence of an ester in an amount of less than 30% by weight of the reaction [mixture] medium.